

Standard Model light-by-light scattering in SANC: analytic and numeric evaluation.

D. Bardin, L. Kalinovskaya, E. Uglov

*Dzhelepov Laboratory for Nuclear Problems, JINR,
ul. Joliot-Curie 6, RU-141980 Dubna, Russia*

Abstract

In this paper we describe the implementation of the SM process $\gamma\gamma \rightarrow \gamma\gamma$ through a fermion and boson loops into the framework of **SANC** system. The computations of this process takes into account non-zero mass of loop particles. We briefly describe additional precomputation modules used for calculation of massive box diagrams. We present the covariant and helicity amplitudes for this process, some particular cases of D_0 and C_0 Passarino–Veltman functions and also numerical results of corresponding **SANC** module evaluation. Whenever possible, we compare the results with those existing in the literature.

E-mails: bardin@nusun.jinr.ru, kalinov@nusun.jinr.ru, corner@nusun.jinr.ru

1 Introduction

SANC is a computer system for semi-automatic calculations of realistic and pseudo-observables for various processes of elementary particle interactions "from the SM Lagrangian to event distributions" at the one-loop precision level for the present and future colliders — TEVATRON, LHC, electron Linear Colliders (ILC, CLIC), muon factories and others. To learn more about available processes in **SANC** see the description in [1, 2] and look at our home pages at JINR and CERN [3].

Light-by-light scattering is one of the most fundamental processes. It proceeds via one-loop box diagrams containing charged particles. The first results for the QED low energy limit of this process were obtained by Euler [4]. Then Karplus and Neumann [5] found a solution for QED in general but complicated way. The QED cross sections in the high energy limit were calculated by Ahiezer [6]. Nowadays there are computations for $\gamma\gamma \rightarrow \gamma\gamma$ process in the electroweak Standard Model [7, 8, 9, 10] and even for two-loop QCD and QED corrections [11].

In this paper we describe the implementation of the SM process $\gamma\gamma \rightarrow \gamma\gamma$ through fermion and boson loops and corresponding precomputation modules into the framework of **SANC** system. The computations of this process take into account non-zero mass of the loop particles.

One should emphasize that the obtained building blocks and procedures of precomputation for box diagrams in QED and EW (as in $\gamma\gamma \rightarrow \gamma\gamma$) is the first step in the creation of environment for implementation of the similar four-bosons processes in the Standard Model (like $\gamma\gamma \rightarrow ZH$, $\gamma\gamma \rightarrow ZZ$ [10]) and in QCD (like $gg \rightarrow \gamma\gamma$, $gg \rightarrow ZZ$, $gg \rightarrow W^+W^-$ etc.).

The paper is organized as follows:

First we discuss some notations and common expression for cross section in section 2.1.

In section 2.2 we discuss diagrams for $\gamma\gamma \rightarrow \gamma\gamma$ process and covariant amplitude tensor structure.

The helicities amplitudes approach [1, 12] and their expressions for light-by-light scattering in general (massive) and in limiting (massless) cases are listed in section 2.3.

In section 3 we shortly describe precomputation strategy and the place of this process on the **SANC** process tree.

The implementation of analytical results and the **SANC** modules concept we describe in section 3.2.

At last in section 3.3 one can find the numerical result and comparisons with those existing in the literature.

Additionally, in Appendix section we list answers for particular cases of D_0 , C_0 and B_0 Passarino–Veltman (PV) functions [13] (see also the book [14]), which are needed for calculation of light-by-light scattering through massive and massless loop particles. Finally, we present strings and basis for covariant amplitude.

2 Light-by-light scattering process

2.1 Notation, cross section

The 4-momenta of incoming photons are denoted by p_1 and p_2 , of the outgoing ones — by p_3 and p_4 . The amplitudes are calculated for scattering of real photons, that is $p_1^2 = 0$, $p_2^2 = 0$, $p_3^2 = 0$, $p_4^2 = 0$. The 4-momentum conservation law reads:

$$p_1 + p_2 - p_3 - p_4 = 0. \quad (1)$$

The Mandelstam variables are:¹

$$\begin{aligned} s &= -(p_1 + p_2)^2 = -2p_1 \cdot p_2, & t &= -(p_1 - p_3)^2 = 2p_1 \cdot p_3, \\ u &= -(p_1 - p_4)^2 = 2p_1 \cdot p_4, & s + t + u &= 0. \end{aligned} \quad (2)$$

¹Note, that in **SANC** we use Pauli metric.

For the $2 \rightarrow 2$ $\gamma\gamma \rightarrow \gamma\gamma$ process, the cross section has the form:

$$d\sigma_{\gamma\gamma \rightarrow \gamma\gamma} = \frac{1}{j} |\mathcal{A}_{\gamma\gamma \rightarrow \gamma\gamma}|^2 d\Phi^{(2)}, \quad (3)$$

where $j = 4\sqrt{(p_1 p_2)^2}$ is the flux, $\mathcal{A}_{\gamma\gamma \rightarrow \gamma\gamma}$ is the covariant amplitude (CA) of the process, and $d\Phi^{(2)}$ is the two body phase space:

$$d\Phi^{(2)} = (2\pi)^4 \delta(p_1 + p_2 - p_3 - p_4) \frac{d^4 p_3 \delta(p_3^2)}{(2\pi)^3} \frac{d^4 p_4 \delta(p_4^2)}{(2\pi)^3}. \quad (4)$$

For the differential cross section one gets:

$$d\sigma_{\gamma\gamma \rightarrow \gamma\gamma} = \frac{1}{128\pi s} |\mathcal{A}_{\gamma\gamma \rightarrow \gamma\gamma}|^2 d\cos\theta, \quad (5)$$

where $s = 4\omega^2$, ω is the photons energy and θ — the scattering angle in the center of mass system (cms).

2.2 Covariant amplitude

The covariant one-loop amplitude corresponds to a result of the straightforward standard calculation of all diagrams contributing to a given process at Born (tree) and one-loop (1-loop) levels.

The CA is being represented in a certain basis, made of strings of Dirac matrices and/or external momenta (structures), contracted with polarization vectors of vector bosons, $\epsilon(p)$, if any. The amplitude also contains kinematical factors and coupling constants and is parameterized by a certain number of Form Factors (FFs), which are denoted by $\mathcal{F}_i$, in general with an index i labeling the corresponding structure. The number of $\mathcal{F}_i$ is equal to the number of independent structures.

The $\gamma\gamma \rightarrow \gamma\gamma$ process in QFT appears due to non-linear effects of interaction with vacuum, so this process has no Born or tree level. Corresponding diagrams start from the one-loop level and in QED there are box diagrams with four internal fermions of equal mass. The number of not identical diagrams (or topologies) is equal to six. But three of them differ from another only by the orientation of the internal fermionic loop, giving the same contribution or a factor 2 to the amplitude. So, only three topologies (st, su and ut channels) remain which are related by simple permutations of external photons in the diagrams shown in Figure 1: su-channel is obtained from st-channel by $p_3 \leftrightarrow p_4$ rotation and ut-channel — by $p_2 \leftrightarrow p_3$. The sum of these fermionic diagrams is a gauge invariant in each generation of particles.




Figure 1: $\gamma\gamma \rightarrow \gamma\gamma$ process QED diagrams

In the EW boson sector we have three types of diagrams to classify: box topologies, pinch topologies and fish topologies (shown in Figure 2). There are three channels of each topologies (st-, su- and ut-channel as in QED) and we have W^+ , W^- , ϕ^+ , ϕ^- and X^+ , X^- (bosons and ghosts) as internal particles in R_ξ -gauge theory.

As in fermionic part we can choose only positive or negative charge bosons and X^+ , X^- ghosts to appear as loop particles and multiply the result by factor 2 to dismiss the double counting diagrams, which differ from another only by the orientation of the loop charge flow.

So, we have three structures (3 channels) in box type of diagrams, twelve structures (3 channels by 4 corresponding pinches) in pinch type and six structures (3 channels by 2 corresponding combinations of propagators — direct and crossed) in fish type of diagrams — each of ones is a sum of the appropriate sets of loop particles diagrams.




Figure 2: $\gamma\gamma \rightarrow \gamma\gamma$ process EW diagrams

The full CA of given process for off-shell photons ($p_i \epsilon_i \neq 0$) with corresponding combinatorial factors can be written as sum of bosonic part minus fermionic and ghost part:

$$\begin{aligned}
 \mathcal{A}_{\gamma\gamma \rightarrow \gamma\gamma} = & +2 \times \left[\sum \text{(box diagram)} + \frac{1}{2} \times \sum \text{(pinch diagram)} + \frac{1}{4} \times \sum \text{(fish diagram)} \right] \\
 & -2 \times \left[\sum \text{(box diagram)} \right] -2 \times \left[\sum \text{(box diagram)} \right]
 \end{aligned} \tag{6}$$

In terms of Lorenz-structures we have:

$$\mathcal{A}_{\gamma\gamma \rightarrow \gamma\gamma} = \sum_{i=1}^{43} \left[\mathcal{F}_i^{\text{bosons}}(s, t, u) + \mathcal{F}_i^{\text{fermions}}(s, t, u) \right] T_i^{\alpha\beta\mu\nu}. \tag{7}$$

The $\mathcal{F}_i$ are normalized by corresponding factors for fermion and boson parts:

$$\begin{aligned}
 C^{\text{fermions}} &= 8\alpha^2 Q_f^4 N_c, \\
 C^{\text{bosons}} &= 12\alpha^2,
 \end{aligned} \tag{8}$$

where α is the fine structure constant, Q_f is the fraction of charge of loop fermion in units of electron charge e , N_c is the number of colours for given fermion, $T_i^{\alpha\beta\mu\nu}$ are tensors defined with an aid of auxiliary strings τ_j presented in the Appendix section IV. The off-shell process contains 43 basis elements, but for the on-shell real photons we need only first 10 structures.

Thus, one gets a minimal number of tensor structures of the CA. It can be written in an explicit form with an aid of scalar $\mathcal{F}_i$. All masses and other parameters dependences are included into these $\mathcal{F}_i$. It is important to emphasize that each basis elements $T_i^{\alpha\beta\mu\nu}$, $i = 1 \div 43$ of tensor structure with Lorenz indices is four times transversal with respect to each external photon:

$$T_i^{\alpha\beta\mu\nu} p_\alpha = T_i^{\alpha\beta\mu\nu} p_\beta = T_i^{\alpha\beta\mu\nu} p_\mu = T_i^{\alpha\beta\mu\nu} p_\nu = 0. \quad (9)$$

$\mathcal{F}_i$ are scalar coefficients in front of basis structures of the CA — projections of CA to complete basis expressions $T_i^{\alpha\beta\mu\nu}$. They are presented as some combinations of scalar Passarino–Veltman functions A_0 , B_0 , C_0 , D_0 [13] and depended on invariants s, t, u and also on fermions and bosons masses. They do not contain UV poles.

The number of terms in $\mathcal{F}_i$ equals to thousands in the case of non-zero mass of the loop particles, but this number reduces greatly for zero loop fermion mass. Full answer for $\mathcal{F}_i$ one can find in the computer system **SANC**. The client part is available on servers [3].

2.3 Helicity amplitudes

In **SANC** we use helicity amplitudes approach [1, 12].

In the expression for CA as one can see in subsection 2.2 one has tensor structures and a set of scalar $\mathcal{F}_i$. To calculate an observable quantity, such as cross section, one needs to make amplitude square, calculate, in general, products of Dirac spinors and contract Lorenz indices with polarization vector. In the standard approach making amplitude square one gets squares for each diagram and their interferences. This leads to a huge number of terms.

In the helicity amplitudes approach we also derive tensor structure and $\mathcal{F}_i$. But the next step is a projection to helicity basis and as a result one gets a set of non-interfering amplitudes, since all of them are characterized by different set of helicity quantum numbers. In this way we can distinguish calculations of Dirac spinors, if they are needed, and contraction of Lorenz indices from calculations of $\mathcal{F}_i$. We can do this before making square of amplitude. So we get a profit on calculation time (less amount of terms due to zero interference) and also more clear step-by-step control.

For the process $\gamma\gamma \rightarrow \gamma\gamma$ one gets:

$$\begin{aligned} \mathcal{A}_{\gamma\gamma \rightarrow \gamma\gamma} &= \sum_{\text{spins}} \left[C^{\text{bosons}} \times \mathcal{H}_{\text{spins}}^{\text{bosons}} + C^{\text{fermions}} \times \mathcal{H}_{\text{spins}}^{\text{fermions}} \right], \\ |\mathcal{A}_{\gamma\gamma \rightarrow \gamma\gamma}|^2 &= \sum_{\text{spins}} \left[C_{\text{bosons}}^2 |\mathcal{H}_{\text{spins}}^{\text{bosons}}|^2 + C_{\text{fermions}}^2 |\mathcal{H}_{\text{spins}}^{\text{fermions}}|^2 \right. \\ &\quad \left. + C^{\text{bosons}} C^{\text{fermions}} \left(\mathcal{H}_{\text{spins}}^{*\text{bosons}} \times \mathcal{H}_{\text{spins}}^{\text{fermions}} + \mathcal{H}_{\text{spins}}^{\text{bosons}} \times \mathcal{H}_{\text{spins}}^{*\text{fermions}} \right) \right]. \quad (10) \end{aligned}$$

Note, the total number of HAs for this process is equal to 16. This corresponds to different combinations of external particles spin projections. For $\gamma\gamma \rightarrow \gamma\gamma$ processes there are 4 photons with two polarizations '+' and '-', so the total number is $2 \cdot 2 \cdot 2 \cdot 2 = 16$. Helicity amplitudes are scalar expressions with factors equal to C^{fermions} for fermions and C^{bosons} for bosons (8).

For bosons part we have:

$$\begin{aligned} \mathcal{H}_{++++}^{\text{bosons}} = \mathcal{H}_{----}^{\text{bosons}} &= -1 + \frac{u-t}{s} \left[B_0(u; M_W, M_W) - B_0(t; M_W, M_W) \right] \\ &\quad + \left(\frac{4M_W^2}{s} + 2\left(\frac{tu}{s^2} - \frac{4}{3}\right) \right) \left(uC_0(0, 0, u; M_W, M_W, M_W) + tC_0(0, 0, t; M_W, M_W, M_W) \right) \\ &\quad - \left(2M_W^2 s \left(\frac{M_W^2}{s} - \frac{4}{3} \right) + \frac{2s^2}{3} \right) \left(D_0(0, 0, 0, 0, s, t; M_W, M_W, M_W, M_W) \right. \\ &\quad \left. + D_0(0, 0, 0, 0, s, u; M_W, M_W, M_W, M_W) + D_0(0, 0, 0, 0, u, t; M_W, M_W, M_W, M_W) \right) \\ &\quad - tu \left(\frac{4M_W^2}{s} + \frac{tu}{s^2} - \frac{4}{3} \right) D_0(0, 0, 0, 0, u, t; M_W, M_W, M_W, M_W), \end{aligned}$$

$$\begin{aligned}
\mathcal{H}_{++++}^{\text{bosons}} = \mathcal{H}_{+--+}^{\text{bosons}} &= \mathcal{H}_{+--+}^{\text{bosons}} = \mathcal{H}_{-+++}^{\text{bosons}} = \mathcal{H}_{----+}^{\text{bosons}} = \mathcal{H}_{---+-}^{\text{bosons}} = \\
\mathcal{H}_{-+-+}^{\text{bosons}} = \mathcal{H}_{+----}^{\text{bosons}} &= 1 - M_W^2 \left(s^2 + t^2 + u^2 \right) \left(\frac{1}{ut} C_0(0, 0, s; M_W, M_W, M_W) \right. \\
&\quad + \frac{1}{su} C_0(0, 0, t; M_W, M_W, M_W) + \frac{1}{st} C_0(0, 0, u; M_W, M_W, M_W) \Big) - M_W^2 \Big(\\
&\quad (2M_W^2 + \frac{st}{u}) D_0(0, 0, 0, 0, s, t; M_W, M_W, M_W, M_W) \\
&\quad + (2M_W^2 + \frac{su}{t}) D_0(0, 0, 0, 0, s, u; M_W, M_W, M_W, M_W) \\
&\quad + (2M_W^2 + \frac{ut}{s}) D_0(0, 0, 0, 0, u, t; M_W, M_W, M_W, M_W) \Big), \\
\mathcal{H}_{+---+}^{\text{bosons}} = \mathcal{H}_{-++-}^{\text{bosons}} &= -1 + \frac{s-t}{u} \left[B_0(s; M_W, M_W) - B_0(t; M_W, M_W) \right] \\
&\quad + \left(\frac{4M_W^2}{u} + 2 \left(\frac{ts}{u^2} - \frac{4}{3} \right) \right) \left(s C_0(0, 0, s; M_W, M_W, M_W) + t C_0(0, 0, t; M_W, M_W, M_W) \right) \\
&\quad - \left(2M_W^2 u \left(\frac{M_W^2}{u} - \frac{4}{3} \right) + \frac{2u^2}{3} \right) \left(D_0(0, 0, 0, 0, u, t; M_W, M_W, M_W, M_W) \right. \\
&\quad + D_0(0, 0, 0, 0, s, u; M_W, M_W, M_W, M_W) + D_0(0, 0, 0, 0, s, t; M_W, M_W, M_W, M_W) \Big) \\
&\quad - ts \left(\frac{4M_W^2}{u} + \frac{ts}{u^2} - \frac{4}{3} \right) D_0(0, 0, 0, 0, s, t; M_W, M_W, M_W, M_W), \\
\mathcal{H}_{+--+}^{\text{bosons}} = \mathcal{H}_{-+-+}^{\text{bosons}} &= -1 + \frac{u-s}{t} \left[B_0(u; M_W, M_W) - B_0(s; M_W, M_W) \right] \\
&\quad + \left(\frac{4M_W^2}{t} + 2 \left(\frac{su}{t^2} - \frac{4}{3} \right) \right) \left(u C_0(0, 0, u; M_W, M_W, M_W) + s C_0(0, 0, s; M_W, M_W, M_W) \right) \\
&\quad - \left(2M_W^2 t \left(\frac{M_W^2}{t} - \frac{4}{3} \right) + \frac{2t^2}{3} \right) \left(D_0(0, 0, 0, 0, s, t; M_W, M_W, M_W, M_W) \right. \\
&\quad + D_0(0, 0, 0, 0, u, t; M_W, M_W, M_W, M_W) + D_0(0, 0, 0, 0, s, u; M_W, M_W, M_W, M_W) \Big) \\
&\quad - su \left(\frac{4M_W^2}{t} + \frac{su}{t^2} - \frac{4}{3} \right) D_0(0, 0, 0, 0, s, u; M_W, M_W, M_W, M_W), \\
\mathcal{H}_{++--}^{\text{bosons}} = \mathcal{H}_{--++}^{\text{bosons}} &= 1 - 2M_W^4 \left(D_0(0, 0, 0, 0, u, t; M_W, M_W, M_W, M_W) + \right. \\
&\quad \left. D_0(0, 0, 0, 0, s, u; M_W, M_W, M_W, M_W) + D_0(0, 0, 0, 0, s, t; M_W, M_W, M_W, M_W) \right) \Big) \Big) \quad (11)
\end{aligned}$$

The analytical answers for scalar Passarino–Veltman functions A_0 , B_0 , C_0 , D_0 [13] are presented in corresponding Appendix sections.

For fermions part we have similar to bosons result with corresponding loop particles masses, but with opposite sign and there is a difference in helicities conserving part of amplitude ($++++$ and $+--+$ permutations), the non-conserving helicities ($++--$) are the same:

$$\begin{aligned}
\mathcal{H}_{++++}^{\text{fermions}} = \mathcal{H}_{----}^{\text{fermions}} &= 1 - \frac{u-t}{s} \left[B_0(u; m_f, m_f) - B_0(t; m_f, m_f) \right] \\
&\quad - \left(\frac{4m_f^2}{s} + 2 \left(\frac{tu}{s^2} - \frac{1}{2} \right) \right) \left(u C_0(0, 0, u; m_f, m_f, m_f) + t C_0(0, 0, t; m_f, m_f, m_f) \right) \\
&\quad + 2m_f^2 s \left(\frac{m_f^2}{s} - \frac{1}{2} \right) \left(D_0(0, 0, 0, 0, s, t; m_f, m_f, m_f, m_f) \right. \\
&\quad + D_0(0, 0, 0, 0, s, u; m_f, m_f, m_f, m_f) + D_0(0, 0, 0, 0, u, t; m_f, m_f, m_f, m_f) \Big) \\
&\quad + tu \left(\frac{4m_f^2}{s} + \frac{tu}{s^2} - \frac{1}{2} \right) D_0(0, 0, 0, 0, u, t; m_f, m_f, m_f, m_f),
\end{aligned}$$

$$\begin{aligned}
\mathcal{H}_{++++}^{\text{fermions}} = \mathcal{H}_{+---}^{\text{fermions}} &= \mathcal{H}_{+--+}^{\text{fermions}} = \mathcal{H}_{-+++}^{\text{fermions}} = \mathcal{H}_{----+}^{\text{fermions}} = \mathcal{H}_{--+--}^{\text{fermions}} = \\
\mathcal{H}_{-+--}^{\text{fermions}} = \mathcal{H}_{+----}^{\text{fermions}} &= -1 + m_f^2 \left(s^2 + t^2 + u^2 \right) \left(\frac{1}{ut} C_0(0, 0, s; m_f, m_f, m_f) \right. \\
&\quad + \frac{1}{su} C_0(0, 0, t; m_f, m_f, m_f) + \frac{1}{st} C_0(0, 0, u; m_f, m_f, m_f) \Big) + m_f^2 \Big(\\
&\quad (2m_f^2 + \frac{st}{u}) D_0(0, 0, 0, 0, s, t; m_f, m_f, m_f, m_f) \\
&\quad + (2m_f^2 + \frac{su}{t}) D_0(0, 0, 0, 0, s, u; m_f, m_f, m_f, m_f) \\
&\quad + (2m_f^2 + \frac{ut}{s}) D_0(0, 0, 0, 0, u, t; m_f, m_f, m_f, m_f) \Big), \\
\mathcal{H}_{+--+}^{\text{fermions}} = \mathcal{H}_{-++-}^{\text{fermions}} &= 1 - \frac{s-t}{u} \left[B_0(s; m_f, m_f) - B_0(t; m_f, m_f) \right] \\
&\quad - \left(\frac{4m_f^2}{u} + 2 \left(\frac{ts}{u^2} - \frac{1}{2} \right) \right) \left(s C_0(0, 0, s; m_f, m_f, m_f) + t C_0(0, 0, t; m_f, m_f, m_f) \right) \\
&\quad + 2m_f^2 u \left(\frac{m_f^2}{u} - \frac{1}{2} \right) \left(D_0(0, 0, 0, 0, u, t; m_f, m_f, m_f, m_f) \right. \\
&\quad + D_0(0, 0, 0, 0, s, u; m_f, m_f, m_f, m_f) + D_0(0, 0, 0, 0, s, t; m_f, m_f, m_f, m_f) \Big) \\
&\quad + ts \left(\frac{4m_f^2}{u} + \frac{ts}{u^2} - \frac{1}{2} \right) D_0(0, 0, 0, 0, s, t; m_f, m_f, m_f, m_f), \\
\mathcal{H}_{-+-+}^{\text{fermions}} = \mathcal{H}_{-++-}^{\text{fermions}} &= 1 - \frac{u-s}{t} \left[B_0(u; m_f, m_f) - B_0(s; m_f, m_f) \right] \\
&\quad - \left(\frac{4m_f^2}{t} + 2 \left(\frac{su}{t^2} - \frac{1}{2} \right) \right) \left(u C_0(0, 0, u; m_f, m_f, m_f) + s C_0(0, 0, s; m_f, m_f, m_f) \right) \\
&\quad + 2m_f^2 t \left(\frac{m_f^2}{t} - \frac{1}{2} \right) \left(D_0(0, 0, 0, 0, s, t; m_f, m_f, m_f, m_f) \right. \\
&\quad + D_0(0, 0, 0, 0, u, t; m_f, m_f, m_f, m_f) + D_0(0, 0, 0, 0, s, u; m_f, m_f, m_f, m_f) \Big) \\
&\quad + su \left(\frac{4m_f^2}{t} + \frac{su}{t^2} - \frac{1}{2} \right) D_0(0, 0, 0, 0, s, u; m_f, m_f, m_f, m_f), \\
\mathcal{H}_{++--}^{\text{fermions}} = \mathcal{H}_{--++}^{\text{fermions}} &= -1 + 2m_f^4 \left(D_0(0, 0, 0, 0, u, t; m_f, m_f, m_f, m_f) + \right. \\
&\quad \left. D_0(0, 0, 0, 0, s, u; m_f, m_f, m_f, m_f) + D_0(0, 0, 0, 0, s, t; m_f, m_f, m_f, m_f) \right). \quad (12)
\end{aligned}$$

In massive case we observe five independent HAs, while in the case of zero loop fermion mass ($m_f = 0$) one gets only four independent HAs which are very compact:

$$\begin{aligned}
\mathcal{H}_{++++}^{\text{fermions}} = \mathcal{H}_{----}^{\text{fermions}} &= -1 + \left(\frac{t-u}{s} \right) (l_u - l_t) - \left(\frac{1}{2} - \frac{ut}{s^2} \right) \left((l_u - l_t)^2 + \pi^2 \right), \\
\mathcal{H}_{+--+}^{\text{fermions}} = \mathcal{H}_{-+++}^{\text{fermions}} &= -1 - i\pi \left(\frac{t-s}{u} \right) - \left[(1+i\pi) \left(\frac{t-s}{u} \right) + 2i\pi \left(\frac{t}{u} \right)^2 \right] l_t - \left(\frac{1}{2} - \frac{st}{u^2} \right) l_t^2, \\
\mathcal{H}_{-+-+}^{\text{fermions}} = \mathcal{H}_{-++-}^{\text{fermions}} &= -1 - i\pi \left(\frac{u-s}{t} \right) - \left[(1+i\pi) \left(\frac{u-s}{t} \right) + 2i\pi \left(\frac{u}{t} \right)^2 \right] l_u - \left(\frac{1}{2} - \frac{su}{t^2} \right) l_u^2. \quad (13)
\end{aligned}$$

where

$$l_t = \ln \left(-\frac{t}{s} \right), \quad l_u = \ln \left(-\frac{u}{s} \right). \quad (14)$$

All the others fermionic HAs are equal to +1.

There are relations among helicity HAs due to C,P,T-invariance. Moreover, there is another relation due to crossing symmetry:

$$\mathcal{H}_{+--+}(s, t, u) = \mathcal{H}_{-+-+}(s, u, t) , \quad (15)$$

but this fact does not mean reducing the number of independent HAs.

3 Process $\gamma\gamma \rightarrow \gamma\gamma$ in SANC

3.1 SANC process tree

In this section we briefly describe analytic modules relevant for $\gamma\gamma \rightarrow \gamma\gamma$.

For boxes the SANC idea of precomputation becomes vitally important [1]. Calculation of some boxes for some particular processes takes so much time that an external user should refrain from repeating pre-computation in the SANC system. Furthermore, the richness of boxes requires a classification. Depending on the type of external lines (f for fermions and b for bosons), we distinguish three large classes of boxes: $ffff$, $ffbb$ and $bbbb$.




Figure 3: $\gamma\gamma \rightarrow \gamma\gamma$ in the SANC process tree

The sum of contributions of fermionic loop boxes form a gauge-invariant and UV-finite subset, which is a consequence of Ward Identity Eq. (9). It is true for sum of bosonic contribution too. So we can distinguish QED and EW part of the process in the analytical calculations.

The precomputation file `AAAA_Box` (see the `SANC` process tree in Figure 3 from `SANC` client [3]) contains the sequence of procedures for calculation of the covariant amplitude. At this step we suppose, that all momenta are incoming (denoted by p 's) and photons are not on-mass-shell. Therefore, these results can be used for other processes which need these parts as building blocks.

When we implement the process $\gamma\gamma \rightarrow \gamma\gamma$ (see `4A_QED` or `EW Processes` branch), we use this building block several times by replacing incoming momenta p 's by corresponding kinematical momenta with the right signs, and calculate $\mathcal{F}_i$ by the module `AA->AA (FF)`, then helicity amplitudes by the module `AA->AA (HA)` and finally — the analytic expression for differential and total process cross section (`XS`) by the module `AA->AA (XS)` for the QED process. For the EW part we use numerical evaluation to get `XS`.

3.2 Stand-alone SANC module

Now let us introduce the `SANC` modules packages concept.

In `SANC` system one has an opportunity of exporting the analytical results for numerical evolution [1]. Moreover, there are tools for checking the implementation of these Standard `SANC FORTRAN` modules (`SSFM`) [2] — the integrator of the process, based on the Vegas algorithm [16].

For light-by-light scattering process the `SSFM` are included in `sanc.4A_v1.00` package, which is available for download from [3]. The numerical results are cross-section distributions from next section. Here we present the technical description of this package — some main flags and the options.

SANC Modules.

SANC NC package (Neutral Current processes modules).

The previous versions of SANC NC package you can find in [SANC Archives](#).

29/07/2009 SANC NC v1.21 package (356 Kb tgz-file) [\[last stable version\]](#)

In v1.21 package some bugs in the EW part are fixed. Details are in the file `CHANGES`.

13/02/2008 SANC NC v1.10 package (300 Kb tgz-file) [\[stable version\]](#)

In v1.10 package the `f1f1HZ` processes are added. For description of the `f1f1HZ` processes please refer to the paper D. Bardin et al., *Comput. Phys. Comm.* 177 (2007) 738-756. Details are in the file `CHANGES`.

This package is intended for calculation of the 1-loop radiative correction to Drell-Yan Neutral Current processes at partonic level. For a Technical Description of this module please refer to the paper *Eur. Phys. J. C* 54 (2008) 451.

SANC CC package (Charged Current processes modules).

The previous versions of SANC CC package you can find in [SANC Archives](#).

16/06/2009 SANC CC v1.30 package (236 Kb tgz-file) [\[last stable version\]](#)

In v1.30 package the modules for new processes are added. Details are in the file `CHANGES`.

05/12/2008 SANC CC v1.20 package (160 Kb tgz-file) [\[stable version\]](#)

In v1.20 package the modules for new processes are added. Details are in the file `CHANGES`.

28/10/2008 SANC CC v1.11 package (131 Kb tgz-file) [\[stable version\]](#)

In v1.11 package some bugs in the QCD soft-virtual part are fixed. Details are in the file `CHANGES`.

This package is intended for calculation of the 1-loop radiative correction to Drell-Yan Charged Current processes at partonic level, see A. Arbuzov et al., *Eur. Phys. J. C* 46 (2006) 407.

SANC 4A package.

19/11/2009 SANC 4A v1.00 package (21 Kb tgz-file) [\[last stable version\]](#)

This package is intended for calculation of the 1-loop full EW correction to light-by-light scattering processes, see D. Bardin et al., "Light-by-light scattering in SANC", *hep-ph/0611188*.

Figure 4: `SANC` modules download web site [3]

The package files:

- `bbbb_main_11.11.F` — the main file;
- `bbbb_ha_11.11.F` — the HA from SANC system;
- `*_r16.f` — the library of special functions and algorithms;
- `*_input.h` — the set of various setups of input parameters;
- README, INSTALL and other instructions files.

In README and INSTALL files one can find instructions how to use the package. The main options one can change in `bbbb_main_11.11.F` — the main file.

There are some main flags and the options with next variants.

- **iha(I)** — choice of helicities sum in cross-section:
I=0 total helicities sum
I=1 ++ helicities sum
I=2 +- helicities sum
- **iqed(I)** — choice of calculations for QED corrections:
I=0 without QED corrections
I=1 with QED corrections
- **iew(I)** — choice of calculations for EW corrections:
I=0 without EW corrections
I=1 with EW corrections
- **gfscheme(I)** — choice of the EW scheme:
I=0 α_0 calculation scheme
I=1 G_F scheme
I=2 G'_F scheme, a test option, when α_0 is replaced by $\alpha_{G_F} = \sqrt{2}G_F M_W^2 (1 - M_W^2/M_Z^2) / \pi$
- **isetaup(I)** — choice of the setup:
I=0 Standard SANC input
I=1 Les Houches Workshop (2005)
I=2 Tevatron-for-LHC Workshop (2006)
I=3 Custom setup (17)
- **start(I)** — choice of start point for 4 order logarithmic scale of $\sqrt{s}$:
I=1q-5 logarithmic scale $\sqrt{s}$ from 0.1 MeV to 1 GeV
I=1q-1 logarithmic scale $\sqrt{s}$ from 1 GeV to 10^4 GeV

To get full EW answer with interference one should set **iqed=1** and **iew=1**.

3.3 Results and comparison

To test our analytical results we calculate final answer for the total cross section for QED part in the massless limit after substitutions of helicity amplitudes and angular integration:

$$\sigma_{\gamma\gamma\rightarrow\gamma\gamma}^{\text{QED}} = \frac{e^8}{2\pi\omega^2} \left(\frac{108}{5} + \frac{13}{2}\pi^2 - 8\pi^2\zeta(3) + \frac{148}{225}\pi^4 - 24\zeta(5) \right). \quad (16)$$

This result was compared with [6] and the complete agreement was found. Also the limit of helicity amplitudes QED was compared separately with [11] and again full agreement was observed. The massive expression of HA for QED and EW parts were compared with [8] and [9].




Figure 5: $\gamma\gamma \rightarrow \gamma\gamma$ SANC and [8] cross-section ($++$ and $+-$) comparison




Figure 6: $\gamma\gamma \rightarrow \gamma\gamma$ SANC total cross-section

The numerical evaluation of analytical expressions for QED, weak and full EW parts within **SSFM** sanc_4A_v1.00 package for cross section were compared with [8] (see Figure 5). The setup is: $\cos \theta < \cos \pi/6$ — cuts; $M_W = 80.22$ GeV; $m_t = 120$ GeV; $\alpha = 1/137.036$; other fermions are massless.

Also according to [9] the sum of helicities cross-sections were calculated for MeV and GeV regions (see Figure 6) with custom setup:

- $\alpha = 1/128$;
 - $\cos \theta < \cos \pi/6$;
 - $M_W = 80.22$ GeV;
 - $m_e = 0.51099892$ MeV, $m_\mu = 0.105658369$ GeV, $m_\tau = 1.77699$ GeV;
 - $m_u = 0.062$ GeV, $m_c = 1.50$ GeV, $m_t = 120.0$ GeV;
 - $m_d = 0.083$ GeV, $m_s = 0.215$ GeV, $m_b = 4.7$ GeV.
- (17)

4 Conclusions

Let us summarise the results.

In this paper the helicities amplitudes expressions for light-by-light scattering in general (massive) and in limiting (massless) cases were listed for QED (fermionic) and weak (bosonic) parts.

We shortly described precomputation strategy of the **SANC** system [1] and the place of this process on the **SANC** process tree.

The implementation of analytical results and the **SSFM** concept were described.

Additionally, we calculated and gave the answers for particular cases of D_0 , C_0 and B_0 Passarino–Veltman (PV) functions for massive and massless loop particles, strings and basis for covariant amplitude.

The **SSFM** sanc_4A_v1.00 package was builded and tested. Its numerical results were compared with those existing in the literature. The package is available for download at web pages [3].

Acknowledgments

The authors are grateful to S. Bondarenko, V. Kolesnikov and A. Saprionov for useful discussions of numerical calculations and to A. Arbuzov for providing us with useful references.

Appendix

To obtain the cross section of the $\gamma\gamma \rightarrow \gamma\gamma$ process in an analytic form we have to compute in an explicit form the master integrals, B_0 , C_0 , D_0 — scalar PV functions [13], [14] — for a particular set of parameters. In D_0 and C_0 functions one can see collinear divergences, but the differential cross section is free of mass singularities which completely cancel in the sum of all terms. The A_0 and B_0 functions contain UV divergences, which cancel in the sum of box contributions. In the process of computation we face also a problem of the “angular edge” divergences, but they are not physical and cancel completely.

I B_0 function

The B_0 function for $\gamma\gamma \rightarrow \gamma\gamma$ process reads:

$$B_0(Q^2; M, M) = \frac{1}{\epsilon} + 2 - \ln\left(\frac{M^2}{\mu^2}\right) - \beta \ln\left(\frac{\beta+1}{\beta-1}\right), \quad (18)$$

where

$$\beta^2 = 1 - \frac{4\widetilde{M}^2}{Q^2},$$

and

$$\widetilde{M}^2 = M^2 - i\epsilon. \quad (19)$$

In zero limit of fermion mass: $M \rightarrow 0$, one gets:

$$\begin{aligned} B_0(s; M, M) &= \frac{1}{\epsilon} + 2 - \left[\ln\left(\frac{s}{\mu^2}\right) - i\pi \right], \\ B_0(t; M, M) &= \frac{1}{\epsilon} + 2 - \left[l_t + \ln\left(\frac{s}{\mu^2}\right) \right], \\ B_0(u; M, M) &= \frac{1}{\epsilon} + 2 - \left[l_u + \ln\left(\frac{s}{\mu^2}\right) \right]. \end{aligned} \quad (20)$$

II C_0 function

The C_0 function is:

$$C_0(0, 0, Q^2; M, M, M) = \int_0^1 dx \int_0^x dy \left(Q^2 y - Q^2 xy + \widetilde{M}^2 \right)^{-1}, \quad (21)$$

After calculations:

$$C_0(0, 0, Q^2; M, M, M) = -\frac{1}{Q^2} \left[\text{Li}_2\left(\frac{1}{x_1}\right) + \text{Li}_2\left(\frac{1}{x_2}\right) \right], \quad (22)$$

where

$$x_{1,2} = \frac{1}{2} (1 \pm \beta). \quad (23)$$

For $M \rightarrow 0$:

$$\begin{aligned}
C_0(0, 0, s; M, M, M) &= -\frac{1}{2s} \left[\ln \left(\frac{M^2}{s} \right) + i\pi \right]^2, \\
C_0(0, 0, t; M, M, M) &= -\frac{1}{2t} \left[\ln \left(\frac{M^2}{s} \right) - l_t \right]^2, \\
C_0(0, 0, u; M, M, M) &= -\frac{1}{2u} \left[\ln \left(\frac{M^2}{s} \right) - l_u \right]^2.
\end{aligned} \tag{24}$$

III D_0 function

The D_0 function looks like:

$$D_0(0, 0, 0, 0, Q^2, P^2; M, M, M, M) = \int_0^1 dx \int_0^x dy \int_0^y dz \left((Q^2 + P^2) xz - P^2 xy - Q^2 yz + P^2 y - P^2 z + \widetilde{M}^2 \right)^{-2} \tag{25}$$

After lengthy calculations:

$$\begin{aligned}
D_0(0, 0, 0, 0, Q^2, P^2; M, M, M, M) &= -\frac{2}{stA_3} \left\{ \right. \\
&\text{Li}_2 \left(\frac{1 + A_3}{A_1 + A_3} \right) - \text{Li}_2 \left(\frac{A_{33}}{A_{13}} \right) + \text{Li}_2 \left(\frac{1 + A_3}{A_2 + A_3} \right) - \text{Li}_2 \left(\frac{A_{33}}{A_{23}} \right) \\
&- \text{Li}_2 \left(-\frac{A_{33}}{A_1 + A_3} \right) - \text{Li}_2 \left(-\frac{A_{33}}{A_2 + A_3} \right) - \text{Li}_2 \left(-\frac{A_{13}}{1 + A_3} \right) - \text{Li}_2 \left(-\frac{A_{23}}{1 + A_3} \right) \\
&\left. - \frac{1}{2} \ln^2 \left(\frac{A_{13}}{1 + A_3} \right) - \frac{1}{2} \ln^2 \left(\frac{A_{23}}{1 + A_3} \right) - 2\zeta(2) + 2i\pi \ln \left(1 + \frac{1}{A_3} \right) \theta(-\text{Im}(A_1 + A_3)) \right\}, \tag{26}
\end{aligned}$$

where

$$\begin{aligned}
A_1 &= \sqrt{1 - \frac{4\widetilde{M}^2}{Q^2}}, \quad A_2 = \sqrt{1 - \frac{4\widetilde{M}^2}{P^2}}, \quad A_3 = \sqrt{1 - \frac{4\widetilde{M}^2(Q^2 + P^2)}{Q^2 P^2}}, \\
A_{33} &= \frac{4\widetilde{M}^2(Q^2 + P^2)}{Q^2 P^2(1 + A_3)}, \quad A_{23} = \frac{4\widetilde{M}^2}{Q^2(A_2 + A_3)}, \quad A_{13} = \frac{4\widetilde{M}^2}{P^2(A_1 + A_3)}.
\end{aligned} \tag{27}$$

$\theta(x)$ is a function of Heaviside.

For $M \rightarrow 0$:

$$\begin{aligned}
D_0(0, 0, 0, 0, s, t; M, M, M, M) &= \frac{2}{st} \left[\ln^2 \left(-\frac{M^2}{t} \right) + \ln \left(-\frac{M^2}{t} \right) l_t - \frac{\pi^2}{2} + i\pi \ln \left(-\frac{M^2}{t} \right) \right] \\
D_0(0, 0, 0, 0, u, t; M, M, M, M) &= \frac{2}{ut} \left[\ln^2 \left(\frac{M^2}{s} \right) - \ln \left(\frac{M^2}{s} \right) (l_t + l_u) - \frac{\pi^2}{2} + l_t l_u \right], \\
D_0(0, 0, 0, 0, s, u; M, M, M, M) &= \frac{2}{su} \left[\ln^2 \left(-\frac{M^2}{u} \right) + \ln \left(-\frac{M^2}{u} \right) l_u - \frac{\pi^2}{2} + i\pi \ln \left(-\frac{M^2}{u} \right) \right].
\end{aligned} \tag{28}$$

IV Strings and basis

To obtain a compact form of structures of the amplitude we choose 14 auxiliary tensorial strings:

$$\begin{aligned}
\tau_1^{\alpha\beta} &= p_{1\beta}p_{2\alpha} + \frac{1}{2}s\delta_{\alpha\beta}, & \tau_2^{\mu\nu} &= p_{3\mu}p_{4\nu} + \frac{1}{2}s\delta_{\mu\nu}, & \tau_3^{\beta\nu} &= p_{2\nu}p_{3\beta} + \frac{1}{2}t\delta_{\beta\nu}, \\
\tau_4^{\alpha\mu} &= p_{1\mu}p_{4\alpha} + \frac{1}{2}t\delta_{\alpha\mu}, & \tau_5^{\alpha\nu} &= p_{1\nu}p_{3\alpha} + \frac{1}{2}u\delta_{\alpha\nu}, & \tau_6^{\beta\mu} &= p_{4\beta}p_{2\mu} + \frac{1}{2}u\delta_{\beta\mu}, \\
\tau_7^\mu &= p_{1\mu} - tu^{-1}p_{2\mu}, & \tau_8^\nu &= p_{1\nu} - ut^{-1}p_{2\nu}, & \tau_9^\beta &= p_{1\beta} - st^{-1}p_{3\beta}, \\
\tau_{10}^\alpha &= p_{2\alpha} - su^{-1}p_{3\alpha}, & \tau_{11}^\mu &= p_{4\mu}, & \tau_{12}^\nu &= p_{3\nu}, & \tau_{13}^\beta &= p_{2\beta}, & \tau_{14}^\alpha &= p_{1\alpha}.
\end{aligned} \tag{29}$$

The complete basis $T_i^{\alpha\beta\mu\nu}$ can be presented in a compact form with an aid of the auxiliary strings τ_j :

$$\begin{aligned}
T_1^{\alpha\beta\mu\nu} &= \tau_1^{\alpha\beta}\tau_2^{\mu\nu}, & T_2^{\alpha\beta\mu\nu} &= \tau_3^{\beta\nu}\tau_4^{\alpha\mu}, & T_3^{\alpha\beta\mu\nu} &= \tau_5^{\alpha\nu}\tau_6^{\beta\mu}, & T_4^{\alpha\beta\mu\nu} &= \tau_1^{\alpha\beta}\tau_7^\mu\tau_8^\nu, \\
T_5^{\alpha\beta\mu\nu} &= \tau_2^{\mu\nu}\tau_9^\beta\tau_{10}^\alpha, & T_6^{\alpha\beta\mu\nu} &= \tau_3^{\beta\nu}\tau_7^\mu\tau_{10}^\alpha, & T_7^{\alpha\beta\mu\nu} &= \tau_4^{\alpha\mu}\tau_8^\nu\tau_9^\beta, & T_8^{\alpha\beta\mu\nu} &= \tau_5^{\alpha\nu}\tau_7^\mu\tau_9^\beta, \\
T_9^{\alpha\beta\mu\nu} &= \tau_6^{\beta\mu}\tau_8^\nu\tau_{10}^\alpha, & T_{10}^{\alpha\beta\mu\nu} &= \tau_7^\mu\tau_8^\nu\tau_9^\beta\tau_{10}^\alpha, & T_{11}^{\alpha\beta\mu\nu} &= \tau_1^{\alpha\beta}\tau_7^\mu\tau_{12}^\nu, & T_{12}^{\alpha\beta\mu\nu} &= \tau_2^{\mu\nu}\tau_9^\beta\tau_{14}^\alpha, \\
T_{13}^{\alpha\beta\mu\nu} &= \tau_3^{\beta\nu}\tau_7^\mu\tau_{14}^\alpha, & T_{14}^{\alpha\beta\mu\nu} &= \tau_4^{\alpha\mu}\tau_8^\nu\tau_{13}^\beta, & T_{15}^{\alpha\beta\mu\nu} &= \tau_5^{\alpha\nu}\tau_7^\mu\tau_{13}^\beta, & T_{16}^{\alpha\beta\mu\nu} &= \tau_6^{\beta\mu}\tau_8^\nu\tau_{14}^\alpha, \\
T_{17}^{\alpha\beta\mu\nu} &= \tau_1^{\alpha\beta}\tau_{11}^\mu\tau_8^\nu, & T_{18}^{\alpha\beta\mu\nu} &= \tau_2^{\mu\nu}\tau_{13}^\beta\tau_{10}^\alpha, & T_{19}^{\alpha\beta\mu\nu} &= \tau_3^{\beta\nu}\tau_{11}^\mu\tau_{10}^\alpha, & T_{20}^{\alpha\beta\mu\nu} &= \tau_4^{\alpha\mu}\tau_{12}^\nu\tau_9^\beta, \\
T_{21}^{\alpha\beta\mu\nu} &= \tau_5^{\alpha\nu}\tau_{11}^\mu\tau_9^\beta, & T_{22}^{\alpha\beta\mu\nu} &= \tau_6^{\beta\mu}\tau_{12}^\nu\tau_{10}^\alpha, & T_{23}^{\alpha\beta\mu\nu} &= \tau_1^{\alpha\beta}\tau_{11}^\mu\tau_{12}^\nu, & T_{24}^{\alpha\beta\mu\nu} &= \tau_2^{\mu\nu}\tau_{13}^\beta\tau_{14}^\alpha, \\
T_{25}^{\alpha\beta\mu\nu} &= \tau_3^{\beta\nu}\tau_{11}^\mu\tau_{14}^\alpha, & T_{26}^{\alpha\beta\mu\nu} &= \tau_4^{\alpha\mu}\tau_{12}^\nu\tau_{13}^\beta, & T_{27}^{\alpha\beta\mu\nu} &= \tau_5^{\alpha\nu}\tau_{11}^\mu\tau_{13}^\beta, & T_{28}^{\alpha\beta\mu\nu} &= \tau_6^{\beta\mu}\tau_{12}^\nu\tau_{14}^\alpha, \\
T_{29}^{\alpha\beta\mu\nu} &= \tau_7^\mu\tau_8^\nu\tau_{13}^\beta\tau_{14}^\alpha, & T_{30}^{\alpha\beta\mu\nu} &= \tau_7^\mu\tau_9^\beta\tau_{12}^\nu\tau_{14}^\alpha, & T_{31}^{\alpha\beta\mu\nu} &= \tau_7^\mu\tau_{10}^\alpha\tau_{12}^\nu\tau_{13}^\beta, & T_{32}^{\alpha\beta\mu\nu} &= \tau_8^\nu\tau_9^\beta\tau_{11}^\mu\tau_{14}^\alpha, \\
T_{33}^{\alpha\beta\mu\nu} &= \tau_8^\nu\tau_{10}^\alpha\tau_{13}^\beta\tau_{11}^\mu, & T_{34}^{\alpha\beta\mu\nu} &= \tau_9^\beta\tau_{10}^\alpha\tau_{11}^\mu\tau_{12}^\nu, & T_{35}^{\alpha\beta\mu\nu} &= \tau_7^\mu\tau_8^\nu\tau_9^\beta\tau_{14}^\alpha, & T_{36}^{\alpha\beta\mu\nu} &= \tau_7^\mu\tau_8^\nu\tau_{10}^\alpha\tau_{13}^\beta, \\
T_{37}^{\alpha\beta\mu\nu} &= \tau_7^\mu\tau_9^\beta\tau_{10}^\alpha\tau_{12}^\nu, & T_{38}^{\alpha\beta\mu\nu} &= \tau_8^\nu\tau_9^\beta\tau_{10}^\alpha\tau_{11}^\mu, & T_{39}^{\alpha\beta\mu\nu} &= \tau_{11}^\mu\tau_{12}^\nu\tau_{13}^\beta\tau_{10}^\alpha, & T_{40}^{\alpha\beta\mu\nu} &= \tau_{11}^\mu\tau_{12}^\nu\tau_{14}^\alpha\tau_9^\beta, \\
T_{41}^{\alpha\beta\mu\nu} &= \tau_{11}^\mu\tau_{13}^\beta\tau_{14}^\alpha\tau_8^\nu, & T_{42}^{\alpha\beta\mu\nu} &= \tau_{12}^\nu\tau_{13}^\beta\tau_{14}^\alpha\tau_7^\mu, & T_{43}^{\alpha\beta\mu\nu} &= \tau_{11}^\mu\tau_{12}^\nu\tau_{13}^\beta\tau_{14}^\alpha.
\end{aligned} \tag{30}$$

References

- [1] A. Andonov *et al.*, CPC 174 (2006), p.481-517.
- [2] A. Andonov *et al.*, CPC 181 (2009), p.305-312.
- [3] WWW: <http://sanc.jinr.ru>, <http://pcphsanc.cern.ch>
- [4] H. Euler, W. Heisenberg, "Quantum Physics" (1936), p.98.
- [5] R. Karplus, M. Neuman, Phys. Rev., v.83 (1951) n.4, p.776.
- [6] A. Ahiezer, V. Beresteky, "Quantum Electrodynamics", p.375, 4 edition, 1981.
- [7] G. Jikia and A. Tkabladze, "Photon-Photon Scattering at the Photon Linear Collider", hep-ph/9312228
- [8] G. Jikia, "Electroweak gauge boson production at $\gamma\gamma$ collider", hep-ph/9710459
- [9] M. Bohm, R. Schuster, Z. Phys. C 63 (1994), p.219.
- [10] Th. Diakonidis *et al.*, "A FORTRAN code for $\gamma\gamma \rightarrow ZZ$ in SM and MSSM", hep-ph/0610085.
- [11] Z. Bern *et al.*, "QCD and QED Corrections to Light-by-Light Scattering", hep-ph/0109079.
- [12] R. Vega, J. Wudka, Phys. Rev. D 53 (1996), p.5286-5292.
- [13] G. Passarino, M. Veltman, Nuclear Physics, B 166 (1979), p.151.
- [14] D. Bardin, G. Passarino, "The Standard Model in the making", Oxford, 1999
- [15] D. Bardin *et al.*, "Light-by-light scattering in SANC", hep-ph/0611188
- [16] G.P. Lepage, J. Comput. Phys. 27 (1978), p.192.